

PubMed

www.pubmed.gov

A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI

[\[Sign In\]](#) [\[Register\]](#)

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search PubMed

for

Go

Clear

Advanced Search

Limits

Preview/Index

History

Clipboard

Details

Display

AbstractPlus

Show

20

Sort By

Send to

All: 1

Review: 0

X

EXHIBIT

9 HUA

DATE

2-18-09

HB

S16

☐ 1: Can J Public Health. 1990 Mar-Apr;81(2):102-6. Links

Comment in:
Can J Public Health. 1990 Sep-Oct;81(5):400-1.

Risk of automobile accidents in cigarette smokers.

Brison RJ.

Division of Emergency Medicine, Queen's University, Kingston, Ontario.

I used a case-control study to identify an increased risk of motor vehicle crash (MVC) in cigarette smokers as compared to non-smokers. Information on smoking status and potential confounding factors was obtained using a self-administered mail-out questionnaire sent to 1,000 persons known to have had a MVC, and a control group of 1,100 persons with no record of MVC in the previous five years. Assessment of relative risk estimates (RR) and adjustment for confounding factors was done by logistic regression analysis. Smokers had a 1.5-fold increase in risk for MVC over non-smokers ($p = .01$). Also, an increased tendency to smoke while driving was associated with greater risk of MVC (X2 trend: $p = .01$). The basis for this association may be: 1. distraction from driving by the act of smoking. 2. behavioural differences between smokers and non-smokers. 3. carbon-monoxide toxicity. Further study is needed to determine the importance of these factors as components of the increase in risk found.

PMID: 2331646 [PubMed - indexed for MEDLINE]





Related Articles

- ▶ Cigarette smoking and acute myocardial infarction. A case-control study from the GIK [Eur J Epidemiol. 1994]
- ▶ Acceptability and concurrent validity of measures to predict older driver involvement [Accid Anal Prev. 2007]
- ▶ Tobacco smoking and periodontal health in a Saudi Arabian population. [Swed Dent J Suppl. 2005]
- ▶ **Review** Smoking and injuries: an overview. [Prev Med. 1994]
- ▶ **Review** Cigarette smoking and risk of non-Hodgkin lymphoma: [Cancer Epidemiol Biomarkers Prev. 2005]

[» See Reviews...](#) [» See All...](#)

Recent Activity

[Turn Off](#) [Clear](#)

-  Risk of automobile accidents in cigarette smokers.
-  Smokers--hazards on road?
-  Can J Public Health. 1990... (2)
-  Can J Public Health. 1990... (0) PubMed

Display

AbstractPlus



Show

20


Sort By

Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)




A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI 
[\[Sign In\]](#) [\[Register\]](#)


All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals

Search PubMed for [Advanced Search](#)


Display Show Sort By

All: 1 Review: 0 

☐ 1: Tob Control. 2005 Jun;14 Suppl 1:i28-32.

 **Full Text**
Tobacco Control

FREE

 **FREE** full text article
in PubMed Central

[Links](#)

Excess injury mortality among smokers: a neglected tobacco hazard.

Wen CP, Tsai SP, Cheng TY, Chan HT, Chung WS, Chen CJ.

Division of Health Policy Research, National Health Research Institutes, Taipei, Taiwan. cwengood@nhri.org.tw

OBJECTIVE: To assess the mortality risks from injuries for smokers and ex-smokers and to quantify the mortality burden of smoking from injury in Taiwan. **METHODS:** Smokers' (and ex-smokers') mortality risks from injuries were compared with that of non-smokers in a merged cohort from Taiwan. A total of 64,319 male subjects were followed up for 12-18 years. Relative risks (RR) (adjusted for age and alcohol use) and 95% confidence intervals (CI) for cause specific injury deaths were calculated using the Cox proportional hazard model. Relative risks of injury mortality were also calculated to assess the presence of dose-response relations with daily smoking quantity. **RESULTS:** Alcohol use adjusted relative mortality risks for all injuries (RR 1.69, 95% CI 1.39 to 2.05) including those from motor vehicle accidents (RR 1.88, 95% CI 1.44 to 2.45) and non-motor vehicle accidents (RR 1.48, 95% CI 1.11 to 1.99) were significantly higher for smokers than non-smokers. Mortality was also increased for most subtypes of non-motor vehicle injuries including falls, fires, and job related injuries. Furthermore, these increases were dose dependent, with the heaviest smokers having the highest risk and the lightest smokers the lowest risk, and ex-smokers, no increase. In 2001, over one fifth (23%) of all male injury deaths in Taiwan was associated with smoking. **CONCLUSION:** This study demonstrated the significant association between fatal injuries and smoking. This relation adds further weight to smoking cessation campaigns.


PMID: 15923444 [PubMed - indexed for MEDLINE]


PMCID: PMC1766175


Related Articles


- ▶ Injury death excesses in smokers: a 1990-95 United States national cohort study. [Inj Prev. 2000]
 - ▶ The mortality risks of smokers in Taiwan: Part I: cause-specific mortality. [Prev Med. 2004]
 - ▶ Impact of smoking on mortality in 80-year-old Japanese from the general popul. [Gerontology. 2008]
 - ▶ **Review** Smoking and injuries: an overview. [Prev Med. 1994]
 - ▶ **Review** When risk factors combine: the interaction between alcohol and smoking for [Addict Behav. 2006]
- » See Reviews... » See All...


Recent Activity

 Excess injury mortality among smokers: a neglected tobacco hazard.

 Smoking and injuries: an overview.



 Risk of automobile accidents in cigarette smokers.

 Smokers--hazards on road?


 [Can J Public Health. 1990... \(2\)](#) PubMed

Display Show Sort By

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)




A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI 
[\[Sign In\]](#) [\[Register\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals

Search PubMed for [Advanced Search](#)

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 1 

☐ 1: Prev Med. 1994 Jul;23(4):515-20.

ELSEVIER [Full-Text Article](#) [Links](#)

Smoking and injuries: an overview.

Sacks JJ, Nelson DE.

Division of Unintentional Injury Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia 30341.

BACKGROUND. Although the disease consequences of cigarette smoking are well documented, smoking may also be associated with increased risk of injury. Our purpose is to provide an overview of this potential association. **METHODS.** We conducted a literature review. **RESULTS.** Cigarettes are the leading cause of death from fire and the second leading cause of fire-related injury. Studies estimate that compared with nonsmokers, smokers appear 1.5 times more likely to have a motor vehicle crash, 1.4-2.5 times more likely to be injured at work, and 2.0 times more likely to suffer other unintentional injuries. A variety of reasons may explain an association between cigarette smoking and injuries; these include (a) direct toxicity; (b) distractibility; (c) smoking-associated medical conditions; and (d) confounding factors, including personality or behavioral characteristics. **CONCLUSIONS.** Smoking may be an independent risk factor for thermal, motor vehicle, occupational, and other unintentional injuries. Nonsmokers may be at increased risk of injury from the presence of smokers in their environments, e.g., from fires. Societal benefits from decreased smoking prevalence are likely to include reduction of both fatal and nonfatal injuries.

PMID: 7971880 [PubMed - indexed for MEDLINE]






Related Articles

- ▶ **Review** Fire injuries, disasters, and costs from cigarettes and cigarette lights: a glol [Prev Med. 2000]
- ▶ Excess injury mortality among smokers: a neglected tobacco hazard. [Tob Control. 2005]
- ▶ Rates of pediatric injuries by 3-month intervals for children 0 to 3 years of age. [Pediatrics. 2003]
- ▶ Injuries as a public health problem in sub-Saharan Africa: epidemiology and prospe [East Afr Med J. 2000]
- ▶ **Review** Intentional and unintentional injuries in women. An overview. [Ann Epidemiol. 1994]

» See Reviews... » See All...


Recent Activity

[Turn Off](#) [Clear](#)

-  Excess injury mortality among smokers: a neglected tobacco hazard.
-  Smoking and injuries: an overview.
-  Risk of automobile accidents in cigarette smokers.
-  Smokers--hazards on road?
-  [Can J Public Health. 1990... \(2\)](#) PubMed

Display AbstractPlus Show 20 Sort By Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



PubMed

A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI

[\[Sign In\]](#) [\[Register\]](#)

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search PubMed

for "Am J Prev Med"[Journal] AND 2006[PDAT] AND

Go

Clear

Advanced Search

Save Search

Limits

Preview/Index

History

Clipboard

Details

Display

AbstractPlus

Show

20

Sort By

Send to

All: 1

Review: 0

We found 1 article in Am J Prev Med 2006:

- ☐ 1: [Am J Prev Med. 2006 Nov;31\(5\):363-8.](#)
- ELSEVIER

FULL-TEXT ARTICLE

Links

Measuring air quality to protect children from secondhand smoke in cars.

Rees VW, Connolly GN.

Harvard School of Public Health, Division of Public Health Practice, Boston, Massachusetts 02115, USA. vrees@hsph.harvard.edu

BACKGROUND: Secondhand tobacco smoke (SHS) is a major, preventable contributor to acute and chronic adverse health outcomes that affect children disproportionately. The predominant source of SHS among children is domestic exposure, and while up to two thirds of U.S. households have car smoking bans, an unacceptable number of children remain vulnerable. To help promote more effective protection through legislation, health communication strategies, or behavioral interventions, data demonstrating the adverse effect of SHS on air quality in cars are needed. METHODS: Secondhand tobacco smoke in a motor vehicle under actual driving conditions was monitored by measuring respirable suspended particles (RSPs) of less than 2.5 microns in diameter, and carbon monoxide. Forty-five driving trials were conducted, using teams of volunteer drivers and smokers recruited from the general community. Three smoking conditions (nonsmoking baseline, active smoking, and immediate post-smoking period, each 5 minutes) were crossed with two ventilation conditions (windows open, closed) in a 3 x 2 within-sessions factorial design. RESULTS: The highest mean observed RSP level was 271 mug/m(3), which is unsafe, particularly for children. Peak RSP levels were considerably higher. RSPs and carbon monoxide increased significantly from baseline after smoking, and these increases were greatest during the closed ventilation condition, compared with open ventilation. CONCLUSIONS: Private passenger cars are a domestic environment with the potential to yield unsafe levels of SHS contaminants. These data may assist policymakers and health advocates to promote protective strategies to ensure smoke-free domestic environments for children.

PMID: 17046406 [PubMed - indexed for MEDLINE]

- Related Articles
- ▶ Air pollution in Boston bars before and after a smoking ban. [BMC Public Health. 2006]

▶ Residual tobacco smoke pollution in used cars for sale: air, dust, and s [Nicotine Tob Res. 2008]

▶ Secondhand smoke and indoor public spaces in Paducah, Kentucky. [J Ky Med Assoc. 2006]

▶ **Review** Flying the smoky skies: secondhand smoke exposure of flight atte [Tob Control. 2004]


▶ **Review** Tobacco industry efforts to present ventilation as an alternative 1 [Tob Control. 2004]
- » See Reviews...


» See All...


Recent Activity


Turn Off


Clear

 Measuring air quality to protect children from secondhand smoke in cars.

 "Am J Prev Med"[Journal] ... (1) PubMed

 Excess injury mortality among smokers: a neglected tobacco hazard.

 Smoking and injuries: an overview.

 Risk of automobile accidents in cigarette smokers.

Display

AbstractPlus

Show

20

Sort By


Send to

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)



PubMed

www.pubmed.gov

A service of the U.S. National Library of Medicine

and the National Institutes of Health

My NCBI

[Sign In] [Register]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search

PubMed

for

"Am J Prev Med"[Journal] AND 2006[PDAT] AND

Go

Clear

Advanced Search

Save Search

Limits

Preview/Index

History

Clipboard

Details

Display

AbstractPlus

Show

20

Sort By

Send to

All: 1

Review: 0

X

We found 1 article in Am J Prev Med 2006:

- ☐ 1: Am J Prev Med. 2006 Nov;31(5):363-8.
- ELSEVIER

FULL-TEXT ARTICLE

Links

Measuring air quality to protect children from secondhand smoke in cars.

Rees VW, Connolly GN.

Harvard School of Public Health, Division of Public Health Practice, Boston, Massachusetts 02115, USA. vrees@hsph.harvard.edu

BACKGROUND: Secondhand tobacco smoke (SHS) is a major, preventable contributor to acute and chronic adverse health outcomes that affect children disproportionately. The predominant source of SHS among children is domestic exposure, and while up to two thirds of U.S. households have car smoking bans, an unacceptable number of children remain vulnerable. To help promote more effective protection through legislation, health communication strategies, or behavioral interventions, data demonstrating the adverse effect of SHS on air quality in cars are needed. METHODS: Secondhand tobacco smoke in a motor vehicle under actual driving conditions was monitored by measuring respirable suspended particles (RSPs) of less than 2.5 microns in diameter, and carbon monoxide. Forty-five driving trials were conducted, using teams of volunteer drivers and smokers recruited from the general community. Three smoking conditions (nonsmoking baseline, active smoking, and immediate post-smoking period, each 5 minutes) were crossed with two ventilation conditions (windows open, closed) in a 3 x 2 within-sessions factorial design. RESULTS: The highest mean observed RSP level was 271 mug/m(3), which is unsafe, particularly for children. Peak RSP levels were considerably higher. RSPs and carbon monoxide increased significantly from baseline after smoking, and these increases were greatest during the closed ventilation condition, compared with open ventilation. CONCLUSIONS: Private passenger cars are a domestic environment with the potential to yield unsafe levels of SHS contaminants. These data may assist policymakers and health advocates to promote protective strategies to ensure smoke-free domestic environments for children.

PMID: 17046406 [PubMed - indexed for MEDLINE]

- Related Articles
- ▶

Air pollution in Boston bars before and after a smoking ban. [BMC Public Health. 2006]
- ▶

Residual tobacco smoke pollution in used cars for sale: air, dust, and s [Nicotine Tob Res. 2008]
- ▶

Secondhand smoke and indoor public spaces in Paducah, Kentucky. [J Ky Med Assoc. 2006]
- ▶

Review Flying the smoky skies: secondhand smoke exposure of flight atte [Tob Control. 2004]
- ▶


Review Tobacco industry efforts to present ventilation as an alternative 1 [Tob Control. 2004]
- » See Reviews...

» See All...


Recent Activity

Turn Off


Clear




Measuring air quality to protect children from secondhand smoke in cars.




"Am J Prev Med"[Journal] ... (1) PubMed



Excess injury mortality among smokers: a neglected tobacco hazard.



Smoking and injuries: an overview.



Risk of automobile accidents in cigarette smokers.

Display

AbstractPlus

Show

20

Sort By



Send to

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer



A service of the U.S. National Library of Medicine
and the National Institutes of Health

My NCBI 2
[Sign In] [Register]

All DatabasesPubMedNucleotideProteinGenomeStructureOMIMPMCJournals

Search PubMed for Go Clear Advanced Search

LimitsPreview/IndexHistoryClipboardDetails

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 0 X

1: Nicotine Tob Res. 2008 Sep;10(9):1467-75.

informaworld Links

Residual tobacco smoke pollution in used cars for sale: air, dust, and surfaces.

Matt GE, Quintana PJ, Hovell ME, Chatfield D, Ma DS, Romero R, Uribe A.
Department of Psychology, San Diego State University, San Diego, CA 92182-4611, USA. gmatt@sciences.sdsu.edu

Regular tobacco use in the enclosed environment of a car raises concerns about longer-term contamination of a car's microenvironment with residual secondhand smoke pollutants. This study (a) developed and compared methods to measure residual contamination of cars with secondhand smoke, (b) examined whether cars of smokers and nonsmokers were contaminated by secondhand smoke, and (c) how smoking behavior and restrictions affected contamination levels. Surface wipe, dust, and air samples were collected in used cars sold by nonsmokers (n = 20) and smokers (n = 87) and analyzed for nicotine. Sellers were interviewed about smoking behavior and restrictions, and car interiors were inspected for signs of tobacco use. Cars of smokers who smoked in their vehicles showed significantly elevated levels of nicotine (p < .001) in dust, on surfaces, and in the air compared with nonsmoker cars with smoking ban. When smokers imposed car smoking bans, air nicotine levels were significantly lower (p < .01), but dust and surface contamination levels remained at similar levels. Smoking more cigarettes in the car and overall higher smoking rate of the seller were significantly associated with higher secondhand smoke contamination of the car (p < .001). Use of a cutpoint for nicotine levels from surface wipe samples correctly identified 82% of smoker cars without smoking bans, 75% of smoker cars with bans, and 100% of nonsmoker cars. Surface nicotine levels provide a relatively inexpensive and accurate method to identify cars and other indoor environments contaminated with residual secondhand smoke. Disclosure requirements and smoke-free certifications could help protect nonsmoking buyers of used cars.

PMID: 19023838 [PubMed - indexed for MEDLINE]

- Related Articles**
- Tobacco use and asking prices of used cars: prevalence, costs, and new opp [Tob Induc Dis. 2008]
 - Measuring air quality to protect children from secondhand smoke in cars. [Am J Prev Med. 2006]
 - Secondhand smoke and indoor public spaces in Paducah, Kentucky. [J Ky Med Assoc. 2006]
 - Review** Flying the smoky skies: secondhand smoke exposure of flight attendants. [Tob Control. 2004]
 - Review** Biomonitoring of inhaled complex mixtures--ambient air, diesel exhaust : [Exp Toxicol Pathol. 2005]
- » See Reviews... » See All...

Recent Activity Turn Off Clear

Residual tobacco smoke pollution in used cars for sale: air, dust, and surfaces.

Measuring air quality to protect children from secondhand smoke in cars.

Q "Am J Prev Med"[Journal] ... (1) PubMed

Excess injury mortality among smokers: a neglected tobacco hazard.

Smoking and injuries: an overview.

Display AbstractPlus Show 20 Sort By Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

NCBI

PubMed

A service of the U.S. National Library of Medicine and the National Institutes of Health

My NCBI [Sign In] [Register]

All DatabasesPubMedNucleotideProteinGenomeStructureOMIMPMCJournals

Search PubMed for "Am J Public Health"[Journal] AND 2008[PDAT] Go Clear Advanced Search Save Search

LimitsPreview/IndexHistoryClipboardDetails

Display AbstractPlus Show 20 Sort By Send to

All: 1 Review: 1

We found 1 article in Am J Public Health 2008:

- ☐ 1: Am J Public Health. 2008 Dec;98(12):2140-5. Epub 2008 Oct 15.
- Final VersionAm J Public HealthLinks

Children's secondhand smoke exposure in private homes and cars: an ethical analysis.

Jarvie JA, Malone RE.

San Francisco Department of Public Health, San Francisco, CA, USA.

Secondhand smoke (SHS) exposure is a known cause of disease among nonsmokers, contributing to lung cancer, heart disease, and sudden infant death syndrome, as well as other diseases. In response to the growing body of scientific literature linking SHS with serious diseases, many countries, states, and cities have established policies mandating smoke-free public spaces. Yet thousands of children remain unprotected from exposure to SHS in private homes and cars. New initiatives targeting SHS in these spaces have raised ethical questions about imposing constraints on private behavior. We reviewed legislation and court cases related to such initiatives and used a principlist approach to analyze the ethical implications of policies banning smoking in private cars and homes in which children are present.

PMID: 18923115 [PubMed - indexed for MEDLINE]

PMCID: PMC2636518 [Available on 2010/12/01]

Related Articles

- ▶ Changing conclusions on secondhand smoke in a sudden infant death syndrom [Pediatrics. 2005]
- ▶ Secondhand smoke in New Zealand homes and cars: exposure, attitudes, and [N Z Med J. 2005]
- ▶ State-specific prevalence of current cigarette smoking ar [MMWR Morb Mortal Wkly Rep. 2001]
- ▶ Review Environmental tobacco smoke: health effects and policie [Annu Rev Public Health. 1997]
- ▶ Review Family and carer smoking control programme [Cochrane Database Syst Rev. 2008]

» See Reviews... | » See All...

Recent Activity

Turn Off Clear

- Children's secondhand smoke exposure in private homes and cars: an ethical
- "Am J Public Health"[Jour... (1)
- Residual tobacco smoke pollution in used cars for sale: air, dust, and surfaces.
- Measuring air quality to protect children from secondhand smoke in cars.
- "Am J Prev Med"[Journal]... (1) PubMed

Display AbstractPlus Show 20 Sort By Send to

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer



NEW WEST
HEALTH SERVICES

Agent Training

Agenda

February 20th, 2009

8:00 a.m. – 3:00 p.m.

8:00 a.m. - 8:45 a.m.	Managing Care in Today's World Dr. Robert Shepard, Medical Director
8:45 a.m. - 9:15 a.m.	New West's Quality Initiatives and Results Monica Wassell, Quality Director
9:15 a.m. - 9:30 a.m.	Break
9:30 a.m. - 10:00 a.m.	How New West Performs UM/UR Cory Hartman, Medical Services Director
10:00 a.m. - 10:30 a.m.	Update on Provider Networks and the Legislative Session Tanya Ask, VP of External & Provider Services
10:30 a.m. - 11:00 a.m.	Medicare Group - A Cost Saving Strategy Bonnie Franklin, Director of Medicare Operations
11:00 a.m. - 11:45 a.m.	Operations and Member Services Dory Hicks, CIO/VP of Operations
11:45 a.m. - 12:45 p.m.	Lunch- New West 2013 David Kibbe, CEO
12:45 p.m. - 1:00 p.m.	Break
1:00 p.m. - 1:30 p.m.	Underwriting Carole Cottrell, Underwriting Manager
1:30 p.m. - 2:15 p. m.	Sales Plan 2009 Greg Loughlin, VP/Strategic Growth
2:15 p.m. - 3:00 p.m.	A Conversation with David Kibbe & Greg Loughlin